V-55 TM IMPROVES NITROGEN METABOLISM

Redox



Dry Soluble Fertilizer 7-25-0

GUARANTEED ANALYSIS

Total NItrogen (N)
7% Ammoniacal Nitrogen
Available Phosphate (P2O5)
Boron (B)
Molybdenum (Mo)
Zinc (Zn)
4% Water soluble Zinc (Zn)
DERIVED FROM ANHYDROUS AMMONIA, PHOSPHORIC ACID, BORIC ACID, SODILIM MOLYRDATE AND ZINC SUITEATE

WHY **V-55**™?

 $V-55^{TM}$ is specifically formulated with phosphorus and micronutrients to improve general nitrogen metabolism and reproductive growth. $V-55^{TM}$ utilizes Redox complexing technology to improve mobility of phosphorus, zinc, and boron in the plant.

PRODUCT USAGE

Apply I-3 lbs./acre as a foliar or soil application every 2 to 4 weeks during the growing season.
Apply I-3 lbs./acre as a foliar or soil application every 2 to 4 weeks during the growing season.



V-55 IMPROVES NITROGEN METABOLISM

V-55™ HANDLING GUIDELINES

PREMIXING

• Premixing is considered a best practice when sprayer agitation is not optimum. Proper hydration is essential for all applications. Recirculate or agitate while adding material.

COMPATIBILITY

- · Always jar test first.
- Redox products are compatible with other Redox products when following product handling guidelines.
- Use caution with reactive materials, such as phosphorus and calcium.
- Avoid extreme shifts in tank pH. When utilizing Redox materials that acidify, check tank pH prior to adding buffers.

TANK MIXING

- Use of an anti-foaming agent is recommended. Fill the tank 50% full with water and initiate tank agitation prior to adding materials.
- Don't add material too quickly—this allows for more thorough hydration.
- The use of inductor assemblies is encouraged.
- · Recirculate or agitate while adding material.
- Total foliar spray content of dry soluble Redox product should not exceed 2% solution (by weight).
- If material is not applied immediately, tank recirculation is required prior to application to ensure uniform product distribution.

REFER TO PRODUCT HANDLING GUIDELINES FOR ADDITIONAL MIXING INSTRUCTIONS.

